


# Dizziness and the Elderly

## FACT SHEET



Author: Lisa Dransfield, PT, DPT, MA



### Consequences of Dizziness in the Elderly


Dizziness, vertigo, and imbalance are not a normal part of aging, but represent the most common complaints that elderly patients bring to their physicians. It is estimated that 2.9% of patients older than 65 years and 3.8% older than 75 years will visit an internist because of dizziness.<sup>1</sup> Unfortunately, dizziness in the elderly can lead to disequilibrium, fear of falling, inactivity, and increased fall risk. In the United States, over 1/3 of older adults fall each year.<sup>2,3</sup> In many older adults, such falls lead to moderate to severe injuries, such as hip fracture, that limit mobility and reduce independent living.<sup>4,5</sup> Dizziness is a major contributing factor to falls in the elderly.<sup>6</sup>

### Causes of Dizziness

Dizziness in the older adult is rarely caused by a single etiology and can be a result of vestibular hypofunction, undiagnosed benign paroxysmal positional vertigo, other medical conditions, polypharmacy, medication side-effects, or a combination of deficiencies in the visual, vestibular, and somatosensory systems. Peripheral vestibular dysfunction is currently thought to account for 48% of dizziness reported by older adults.<sup>1</sup>


### The Role of Physical Therapy in Managing Dizziness

Physical therapists are the practitioners of choice in the rehabilitation and management of vestibular-related balance disorders.<sup>7</sup> A physical therapist specializing in vestibular rehabilitation will perform a thorough examination to determine the cause of symptoms and devise a unique treatment plan to eliminate or minimize dizziness and its consequences. Additional goals of vestibular rehabilitation include reducing fall risk, screening for and correcting benign paroxysmal positional vertigo (often undetected in the elderly<sup>8</sup>), stabilizing balance, retraining the proprioceptive system, improving gaze stability, gait training, and enabling optimal function. Fortunately, aging does not adversely affect rehabilitation outcomes. Research demonstrates that vestibular rehabilitation is just as successful in remediating symptoms in the elderly as in a younger population.<sup>9</sup>



1111 North Fairfax Street  
Alexandria, VA 22314-1488  
Phone: 800-999-2782,  
Ext 3237  
Fax: 703-706-8578  
Email: [neuro@apta.org](mailto:neuro@apta.org)  
[www.neuropt.org](http://www.neuropt.org)

Dizziness and the Elderly



**References:**

1. Herdman SJ. *Vestibular Rehabilitation*. 3<sup>rd</sup> ed. Philadelphia: F.A. Davis Company; 2007.
2. Hornbrook MC, Stevens VJ, Wingfield DJ, Hollis JF, Greenlick MR, Ory MG. Preventing falls among community-dwelling older persons: results from a randomized trial. *Gerontologist*. 1994; 34: 16-23.
3. Hausdorff JM, Rios DA, Edelber HK. Gait variability and fall risk in community-living older adults: a 1-year prospective study. *Arch Phys Med Rehabil*. 2001; 82: 1050-1056.
4. Alexander BH, Rivara FP, Wolf ME. The cost and frequency of hospitalization for fall-related injuries in older adults. *Am J Public Health*. 1992; 82: 1020-1023.
5. Sterling DA, O'Connor JA, Bonadies J. Geriatric falls: injury severity is high and disproportionate to mechanism. *J Trauma*. 2001; 50: 116-119.
6. Agarwal Y, Carey JP, Della Santina CC, Schubert MC, Minor LB. Disorders of balance and vestibular function in US adults: data from the National Health and Nutrition Examination Survey, 2001-2004. *Arch Intern Med*. 2009;169(10):938-944.
7. Physical Therapists as Practitioners of Choice to Rehabilitate Persons with Vestibular-Related Balance Disorders (House of Delegates 05-07-18-18). [http://www.apta.org/AM/Template.cfm?Section=Policies\\_and\\_Bylaws&TEMPLATE=/CM/ontentDisplay.cfm&CONTENTID=67331](http://www.apta.org/AM/Template.cfm?Section=Policies_and_Bylaws&TEMPLATE=/CM/ontentDisplay.cfm&CONTENTID=67331). Accessed July 17, 2010.
8. Oghalai JS, Manolidis S, Barth JL, Stewart MG, Jenkins HA. Unrecognized benign paroxysmal positional vertigo in elderly patients. *Otolaryngol Head Neck Surg*. 2000;122(5):630-634.
9. Whitney SL, Wrisley DM, Marchetti GF, Furman JM. The effect of age on vestibular rehabilitation outcomes. *Laryngoscope*. 2002; 112: 1785-1790.



1111 North Fairfax Street  
Alexandria, VA 22314-1488  
Phone: 800-999-2782,  
Ext 3237  
Fax: 703-706-8578  
Email: [neuropt@apta.org](mailto:neuropt@apta.org)  
[www.neuropt.org](http://www.neuropt.org)