


Medications and Dizziness

FACT SHEET



Author: Melissa Bloom, PT, DPT



Vestibular Suppressants

For the first few days of a vestibular crisis, Meclizine and other vestibular suppressants can be helpful to control severe vertigo, dizziness, and nausea. However, chronic use is counterproductive to the central nervous system's (CNS) compensation process. Vestibular suppressants decrease the CNS's ability to compensate for vestibular loss and can prolong recovery time.¹⁻⁴ Vestibular suppressants have the added risk of producing drowsiness, cognitive deficits, difficulty driving, and increasing the possibility of falls.⁵ The recommendation to decrease and eliminate vestibular suppressants has been well documented in the literature.^{1,3-9}

Vestibular Neuritis

It is thought that while vestibular neuritis is caused by a virus, the symptoms are due to the inflammation process causing compression of the vestibular nerve.¹⁰⁻¹² Studies have shown the effectiveness of corticosteroids to treat the impairments of vestibular neuritis^{11,12} and the benefits of corticosteroids over antiviral or placebo medications.¹¹

Benign Paroxysmal Positional Vertigo (BPPV)

The practice guideline from the American Academy of Otolaryngology - Head and Neck Surgery recommends that clinicians do not treat BPPV with medication and instead advocates the use of canalith repositioning maneuvers. Vestibular suppressants are only recommended for the short term management of severe nausea or vomiting. Clinicians should instead offer education regarding medications, and their ability to cause potential harm versus benefit.⁵ The practice guideline from the American Academy of Neurology states that research regarding the use of medication to treat BPPV is "inadequate or conflicting" and also recommends the use of canalith repositioning maneuvers to treat BPPV.¹³ There has been no strong evidence to support the use of medications in the resolution of BPPV^{5,13} and there is no evidence demonstrating that vestibular suppressants are as effective as canalith repositioning maneuvers.⁵


Migraine

Medications shown to be effective in treating migraine associated dizziness are



1111 North Fairfax Street
Alexandria, VA 22314-1488
Phone: 800-999-2782,
Ext 3237
Fax: 703-706-8578
Email: neuro@apta.org
www.neuropt.org

Medications and Dizziness



antidepressants, beta blockers, calcium channel blockers, and anticonvulsant medications.^{14, 15} Patients who received medicine as treatment for migraine associated dizziness have shown significantly better outcomes than patients who were not medicated.¹⁶

Meniere's Disease

Diuretics can significantly decrease the frequency and severity of Meniere's attacks.^{14,17,18} The combination of medication and reduced dietary sodium is thought to decrease endolymphatic volume. During an acute attack of Meniere's disease, medication for symptom control is recommended.

Anxiety

Anxiety associated with dizziness may or may not be correlated with vestibular dysfunction. Anti-anxiety medications (Benzodiazepines) and anti-depressants (SSRIs) have been recommended for patients with anxiety, dizziness, and balance impairments.¹⁵

In summary, medications can be useful in the management of certain vestibular disorders. However, they are not always indicated or necessary for optimal recovery of function. Careful consideration should be given to the patient's diagnosis and when in the course of recovery medications are administered.

References:


1. Kondrad HR, Tomlison D, Stockwell CW, et al. Rehabilitation therapy for patients with disequilibrium and balance disorders. *Otolaryngol Head Neck Surg* 1992; 107:105-108.
2. Baloh RW. Vertigo. *The Lancet* 1998; 352:1841-1846.
3. Peppard SB. Effect of drug therapy on compensation from vestibular injury. *Laryngoscope* 1996; 8:878-98.
4. Shepard NT, Telian SA. Programmatic vestibular rehabilitation. *Otolaryngol Head Neck Surg* 1995; 112(1): 173-182
5. Bhattacharyya N, Baugh RF, Orvidas L, et al. Clinical practice guideline: Benign paroxysmal positional vertigo. *Otolaryngol Head Neck* 2008; 139:S47-S81.
6. Whitney SL, Rossi MM. Efficacy of vestibular rehabilitation. *Otolaryngol Clin North Am* 2000; 33(3):659-672.
7. Robertson D, Ireland D. Evaluation and treatment of uncompensated unilateral vestibular disease. *Otolaryngol Clin North Am* 1997; 30:745-757.
8. Hanley K, O'Dowd T, Considine N. A systematic review of vertigo in primary care. *Br J Gen Pract* 2001; 51(469):666-671.
9. Brandt T. Management of vestibular disorders. *J Neurol* 2000; 247:491-499.
10. Baloh RW. Vestibular Neuritis. *N Engl J Med* 2003; 348:1027-1032.
11. Strupp M, Zingler VC, Arbusow V, et al. Methylprednisolone, Valacyclovir, or the Combination for Vestibular Neuritis. *N Engl J Med* 2004; 351(4):354-361.



1111 North Fairfax Street
Alexandria, VA 22314-1488
Phone: 800-999-2782,
Ext 3237
Fax: 703-706-8578
Email: neuropt@apta.org
www.neuropt.org

AMERICAN PHYSICAL THERAPY ASSOCIATION, SECTION ON NEUROLOGY

Medications and Dizziness



12. Ariyasu L, Byl FM, Sprague MS, et al. The beneficial effect of methylprednisolone. Arch Otolaryngol Head Neck Surg 1990; 116:700-703.
13. Fife TD, Iverson DJ, Lempert T, et al. Practice parameter: therapies for benign paroxysmal positional vertigo (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology. Neurology 2008; 70(22) 2067-74
14. Chawla N, Olshaker JS. Diagnosis and management of dizziness and vertigo. Med Clin N Am 2006; 90:291-304.
15. Furman JM, Balaban CD, Jacob RG, et al. Migraine-anxiety related dizziness (MARD): a new disorder? J Neurol Neurosurg Psychiatry 2005; 76:1-8.
16. Whitney SL, Wrisley DM, Brown KE, et al. Physical therapy for migraine-related vestibulopathy and vestibular dysfunction with history of migraine. The Laryngoscope; 110 (9):1528-1534.
17. Claes J, Van de Heyning PH. Medical treatment of Meniere's disease: a review of literature. Acta Oto-laryngologica 1997; 117(s526):37-42.
18. Sajjadi H. Medical management of Meniere's disease. Otolaryngol Clin N Am 2002; 35:581-589.



1111 North Fairfax Street
Alexandria, VA 22314-1488
Phone: 800-999-2782,
Ext 3237
Fax: 703-706-8578
Email: neuropt@apta.org
www.neuropt.org